V-Fib/Pulseless V-Tach

Oshkosh Fire Dept

Notes:
- When advanced airway is in place, ventilation shall not be more than 10 breaths per min.

Witness by OFD Personnel
- They go into V-Fib/V-Tach in front of you

Yes

Defibrillate 200 joules

CPR for 2 min. with indicated study device, intubate and establish an IV s/normal saline.
DO NOT stop compressions for procedures

Defibrillate 200 joules

CPR for 2 min. w/indicated study device
Epinephrine 1 mg 1:10,000 IVP/IO every 3-5 min.
OR
Epinephrine 2 mg 1:1,000 ETT (if IV/IO cannot be established)

Defibrillate 200 joules

CPR for 2 min. w/indicated study device
Amiodarone 300 mg IVP/IO

Defibrillate 200 joules

- Continue w/CPR, Defibrillate every 2 min, and Epinephrine every 3-5 min.
- 2 Grams Magnesium Sulfate for Refractory V-Fib/V-Tach and Torsades de pointes

No

CPR for 2 min. with indicated study device, intubate and establish an IV s/normal saline.
DO NOT stop compressions for procedures

Defibrillate 200 joules

CPR for 2 min. w/indicated study device
Epinephrine 1 mg 1:10,000 IVP/IO every 3-5 min.
OR
Epinephrine 2 mg 1:1,000 ETT (if IV/IO cannot be established)

Defibrillate 200 joules

CPR for 2 min. w/indicated study device
Amiodarone 300 mg IVP/IO

Defibrillate 200 joules

- Continue w/CPR, Defibrillate every 2 min, and Epinephrine every 3-5 min.
- 2 Grams Magnesium Sulfate for Refractory V-Fib/V-Tach and Torsades de pointes

Steve Stroman, MD, FACEP, FAAEM, EMT-P
Medical Director, Oshkosh Fire Dept